

Rhode Island Department of Environmental Management Determination of a Groundwater Quality Standard for:

Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonate (PFOS)

Pursuant to Rule 11.2.2 of the Rhode Island Department of Environmental Management (RIDEM) Groundwater Quality Rules, the Director of RIDEM has by means of this document determined that the groundwater quality standard for groundwater classified GAA and GA for PFOA, PFOS or a combination of PFOA and PFOS shall be 70 parts per trillion.

Background:

PFOA and PFOS are part of a class of chemicals referred to as per- and polyfluoroalkyl substances that are used in a variety of products and applications including non-stick cookware, upholstered furniture, clothing, food packaging, and firefighting foam. Facilities that have the potential to contain these chemicals due to use or disposal include industrial factories, airports, fire training academies, landfills, and other waste disposal facilities.

In 2016, the US Environmental Protection Agency (EPA) established a Drinking Water Health Advisory for PFOA, PFOS or a combination of PFOA and PFOS at 70 parts per trillion. Studies indicate that exposure to PFOA and PFOS over the EPA Health Advisory level may result in adverse health effects, including developmental effects to fetuses during pregnancy or to breastfed infants; cancer; and effects to the liver, immune system, or thyroid.

The EPA Health Advisory Fact Sheet is attached and further information on the Health Advisory can be found on EPA's webpage at: https://www.epa.gov/ground-water-and-drinking-water-health-advisories-pfoa-and-pfos

Need for Groundwater Quality Standard:

The establishment of a groundwater quality standard for PFOA and PFOS is justified by:

- Widespread use of these chemicals;
- Potential for these chemicals to be in groundwater;
- The presence of these chemicals in groundwater in other New England states; and
- The detection of these chemicals in groundwater in public water systems in RI.

The Rhode Island Department of Health (RIDOH) is currently sampling selected public wells for these chemicals because of the increasing public health concerns about these chemicals in drinking water supplies. If these chemicals are detected in a groundwater supply, RIDEM will be responsible to investigate the source. Once the source of these chemicals has been identified, RIDEM will oversee actions by the source owner to mitigate any present or past release of PFOA

or PFOS. A groundwater quality standard is necessary to compel action by an identified source owner and to establish a target for any necessary remediation.

Determination of Groundwater Quality Standard for PFOA and PFOS

The groundwater quality standard for groundwater classified GAA and GA in RI is suitability for drinking water use without treatment as specified in RI General Laws, Section 46-13.1. The RIDEM Groundwater Quality Rules (Rule 11), therefore, determined that the standards for groundwater classified GAA and GA would be the federal drinking water maximum contaminant levels established pursuant to the federal Safe Drinking Water Act. For substances that do not have a maximum contaminant level, the GAA and GA standard "... shall be determined by the Director on a case-by-case basis using United States Environmental Protection Agency health advisories and other public health information" (Rule 11.2.2).

Based on the following:

- The US EPA has designated 70 parts per trillion as the Health Advisory for PFOA and PFOS or a combination of PFOA and PFOS;
- The RIDOH will be using the EPA Health Advisory in determining the appropriate response to protect public health when these chemicals are detected in public and private drinking water wells; and
- RIDEM should be consistent with the RI Department of Health drinking water standards and health advisories in adopting groundwater quality standards for groundwater classified GAA and GA, which is to be protected to maintain drinking water quality.

Therefore, RIDEM is adopting 70 parts per trillion as the groundwater quality standard for PFOA, PFOS or a combination of PFOA and PFOS where the groundwater is classified GAA or GA.

Janet Coit

Director

Attachment: EPA Fact Sheet PFOA & PFOS Drinking Water Health Advisories, November 2016



FACT SHEET PFOA & PFOS Drinking Water Health Advisories



Overview

EPA has established health advisories for PFOA and PFOS based on the agency's assessment of the latest peer-reviewed science to provide drinking water system operators, and state, tribal and local officials who have the primary responsibility for overseeing these systems, with information on the health risks of these chemicals, so they can take the appropriate actions to protect their residents. EPA is committed to supporting states and public water systems as they determine the appropriate steps to reduce exposure to PFOA and PFOS in drinking water. As science on health effects of these chemicals evolves, EPA will continue to evaluate new evidence.

Background on PFOA and PFOS

PFOA and PFOS are fluorinated organic chemicals that are part of a larger group of chemicals referred to as perfluoroalkyl substances (PFASs). PFOA and PFOS have been the most extensively produced and studied of these chemicals. They have been used to make carpets, clothing, fabrics for furniture, paper packaging for food and other materials (e.g., cookware) that are resistant to water, grease or stains. They are also used for firefighting at airfields and in a number of industrial processes.

Because these chemicals have been used in an array of consumer products, most people have been exposed to them. Between 2000 and 2002, PFOS was voluntarily phased out of production in the U.S. by its primary manufacturer. In 2006, eight major companies voluntarily agreed to phase out their global production of PFOA and PFOA-related chemicals, although there are a limited number of ongoing uses. Scientists have found PFOA and PFOS in the blood of nearly all the people they tested, but these studies show that the levels of PFOA and PFOS in blood have been decreasing. While consumer products and food are a large source of exposure to these chemicals for most people, drinking water can be an additional source in the small percentage of communities where these chemicals have contaminated water supplies. Such contamination is typically localized and associated with a specific facility, for example, an industrial facility where these chemicals were produced or used to manufacture other products or an airfield at which they were used for firefighting.

EPA's 2016 Lifetime Health Advisories

EPA develops health advisories to provide information on contaminants that can cause human health effects and are known or anticipated to occur in drinking water. EPA's health advisories are non-enforceable and non-regulatory and provide technical information to states agencies and other public health officials on health effects, analytical methodologies, and treatment technologies associated with drinking water contamination. In 2009, EPA published provisional health advisories for PFOA and PFOS based on the evidence available at that time. The science has evolved since then and EPA is now replacing the 2009 provisional advisories with new, lifetime health advisories.

FACT SHEET PFOA & PFOS Drinking Water Health Advisories

EPA's 2016 Lifetime Health Advisories, continued

To provide Americans, including the most sensitive populations, with a margin of protection from a lifetime of exposure to PFOA and PFOS from drinking water, EPA established the health advisory levels at 70 parts per trillion. When both PFOA and PFOS are found in drinking water, the <u>combined</u> concentrations of PFOA and PFOS should be compared with the 70 parts per trillion health advisory level. This health advisory level offers a margin of protection for all Americans throughout their life from adverse health effects resulting from exposure to PFOA and PFOS in drinking water.

How the Health Advisories were developed

EPA's health advisories are based on the best available peer-reviewed studies of the effects of PFOA and PFOS on laboratory animals (rats and mice) and were also informed by epidemiological studies of human populations that have been exposed to PFASs. These studies indicate that exposure to PFOA and PFOS over certain levels may result in adverse health effects, including developmental effects to fetuses during pregnancy or to breastfed infants (e.g., low birth weight, accelerated puberty, skeletal variations), cancer (e.g., testicular, kidney), liver effects (e.g., tissue damage), immune effects (e.g., antibody production and immunity), thyroid effects and other effects (e.g., cholesterol changes).

EPA's health advisory levels were calculated to offer a margin of protection against adverse health effects to the most sensitive populations: fetuses during pregnancy and breastfed infants. The health advisory levels are calculated based on the drinking water intake of lactating women, who drink more water than other people and can pass these chemicals along to nursing infants through breastmilk.

Recommended Actions for Drinking Water Systems

Steps to Assess Contamination

If water sampling results confirm that drinking water contains PFOA and PFOS at individual or combined concentrations greater than 70 parts per trillion, water systems should quickly undertake additional sampling to assess the level, scope and localized source of contamination to inform next steps

Steps to Inform

If water sampling results confirm that drinking water contains PFOA and PFOS at individual or combined concentrations greater than 70 parts per trillion, water systems should promptly notify their State drinking water safety agency (or with EPA in jurisdictions for which EPA is the primary drinking water safety agency) and consult with the relevant agency on the best approach to conduct additional sampling.

Drinking water systems and public health officials should also promptly provide consumers with information about the levels of PFOA and PFOS in their drinking water. This notice should include specific information on the risks to fetuses during pregnancy and breastfed and formula-fed infants from exposure to drinking water with an individual or combined concentration of PFOA and PFOS above EPA's health advisory level of 70 parts per trillion. In addition, the notification should include actions they are taking and identify options that consumers may consider to reduce risk such as seeking an alternative drinking water source, or in the case of parents of formula-fed infants, using formula that does not require adding water.

FACT SHEET PFOA & PFOS Drinking Water Health Advisories

Recommended Actions for Drinking Water Systems, continued

Steps to Limit Exposure

A number of options are available to drinking water systems to lower concentrations of PFOA and PFOS in their drinking water supply. In some cases, drinking water systems can reduce concentrations of perfluoroalkyl substances, including PFOA and PFOS, by closing contaminated wells or changing rates of blending of water sources. Alternatively, public water systems can treat source water with activated carbon or high pressure membrane systems (e.g., reverse osmosis) to remove PFOA and PFOS from drinking water. These treatment systems are used by some public water systems today, but should be carefully designed and maintained to ensure that they are effective for treating PFOA and PFOS. In some communities, entities have provided bottled water to consumers while steps to reduce or remove PFOA or PFOS from drinking water or to establish a new water supply are completed.

Many home drinking water treatment units are certified by independent accredited third party organizations against American National Standards Institute (ANSI) standards to verify their contaminant removal claims. NSF International (NSF®) has developed a protocol for NSF/ANSI Standards 53 and 58 that establishes minimum requirements for materials, design and construction, and performance of point-of-use (POU) activated carbon drinking water treatment systems and reverse osmosis systems that are designed to reduce PFOA and PFOS in public water supplies. The protocol has been established to certify systems (e.g., home treatment systems) that meet the minimum requirements. The systems are evaluated for contaminant reduction by challenging them with an influent of $1.5\pm30\%$ µg/L (total of both PFOA and PFOS) and must reduce this concentration by more than 95% to 0.07 µg/L or less (total of both PFOA and PFOS) throughout the manufacturer's stated life of the treatment system. Product certification to this protocol for testing home treatment systems verifies that devices effectively reduces PFOA and PFOS to acceptable levels.

Other Actions Relating to PFOA and PFOS

Between 2000 and 2002, PFOS was voluntarily phased out of production in the U.S. by its primary manufacturer, 3M. EPA also issued regulations to limit future manufacturing, including importation, of PFOS and its precursors, without first having EPA review the new use. A limited set of existing uses for PFOS (fire resistant aviation hydraulic fluids, photography and film products, photomicrolithography process to produce semiconductors, metal finishing and plating baths, component of an etchant) was excluded from these regulations because these uses were ongoing and alternatives were not available.

In 2006, EPA asked eight major companies to commit to working toward the elimination of their production and use of PFOA, and chemicals that degrade to PFOA, from emissions and products by the end of 2015. All eight companies have indicated that they have phased out PFOA, and chemicals that degrade to PFOA, from emissions and products by the end of 2015. Additionally, PFOA is included in EPA's proposed Toxic Substance Control Act's Significant New Use Rule (SNUR) issued in January 2015 which will ensure that EPA has an opportunity to review any efforts to reintroduce the chemical into the marketplace and take action, as necessary, to address potential concerns.

FACT SHEET PFOA & PFOS Drinking Water Health Advisories

Other Actions Relating to PFOA and PFOS, continued

EPA has not established national primary drinking water regulations for PFOA and PFOS. EPA is evaluating PFOA and PFOS as drinking water contaminants in accordance with the process required by the Safe Drinking Water Act (SDWA). To regulate a contaminant under SDWA, EPA must find that it: (1) may have adverse health effects; (2) occurs frequently (or there is a substantial likelihood that it occurs frequently) at levels of public health concern; and (3) there is a meaningful opportunity for health risk reduction for people served by public water systems.

EPA included PFOA and PFOS among the list of contaminants that water systems are required to monitor under the third Unregulated Contaminant Monitoring Rule (UCMR 3) in 2012. Results of this monitoring effort are updated regularly and can be found on the publicly-available National Contaminant Occurrence Database (NCOD) (https://www.epa.gov/dwucmr/occurrence-data-unregulated-contaminant-monitoringrule#3). In accordance with SDWA, EPA will consider the occurrence data from UCMR 3, along with the peer reviewed health effects assessments supporting the PFOA and PFOS Health Advisories, to make a regulatory determination on whether to initiate the process to develop a national primary drinking water regulation.

In addition, EPA plans to begin a separate effort to determine the range of PFAS for which an Integrated Risk Information System (IRIS) assessment is needed. The IRIS Program identifies and characterizes the health hazards of chemicals found in the environment. IRIS assessments inform the first two steps of the risk assessment process: hazard identification, and dose-response. As indicated in the 2015 IRIS Multi-Year Agenda, the IRIS Program will be working with other EPA offices to determine the range of PFAS compounds and the scope of assessment required to best meet Agency needs. More about this effort can be found at https://www.epa.gov/iris/iris-agenda.

Non-Drinking Water Exposure to PFOA and PFOS

These health advisories only apply to exposure scenarios involving drinking water. They are not appropriate for use, in identifying risk levels for ingestion of food sources, including: fish, meat produced from livestock that consumes contaminated water, or crops irrigated with contaminated water.

The health advisories are based on exposure from drinking water ingestion, not from skin contact or breathing. The advisory values are calculated based on drinking water consumption and household use of drinking water during food preparation (e.g., cooking or to prepare coffee, tea or soup). To develop the advisories, EPA considered non-drinking water sources of exposure to PFOA and PFOS, including: air, food, dust, and consumer products. In January 2016 the Food and Drug Administration amended its regulations to no longer allow PFOA and PFOS to be added in food packaging, which will likely decrease one source of non-drinking water exposure.

Where Can I Learn More?

- EPA's Drinking Water Health Advisories for PFOA and PFOS can be found at: https://www.epa.gov/ground-water-and-drinking-water/drinking-water-health-advisories-pfoa-and-pfos
- PFOA and PFOS data collected under EPA's Unregulated Contaminant Monitoring Rule are available: https://www.epa.gov/dwucmr/occurrence-data-unregulated-con taminant-monitoring-rule
- EPA's stewardship program for PFAS related to TSCA: <a href="https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/and-polyfluoroalkyl-substances-pfass-under-tsca
- EPA's research activities on PFASs can be found at: http://www.epa.gov/chemical-research/
 perfluorinated-chemical-pfc-research
- The Agency for Toxic Substances and Disease Registry's Perflourinated Chemicals and Your Health webpage at: http://www.atsdr.cdc.gov/PFC/

